GRAINS FOR THE DRY LANDS OF CENTRAL OREGON

L. R. BREITHAUPT

Superintendent, Harney Branch Experiment Station
Burns, Oreg.

FARMERS' BULLETIN 800

UNITED STATES DEPARTMENT OF AGRICULTURE

Contribution from the Bureau of Plant Industry
WM. A. TAYLOR, Chief

Washington, D. C. April, 1917

WASHINGTON : GOVERNMENT PRINTING OFFICE : 1917

THIS bulletin discusses the production of small grains on nonirrigated lands in central and southeastern Oregon at elevations between 4,000 and 5,000 feet. The important crops for these dry-farmed lands are winter wheat and rye and spring wheat, rye, oats, and barley. Field peas are also of value as a legume for forage and grain. Dependence upon grain farming alone, however, is not advisable in this region; the grain crops should be grown for the winter feeding and the finishing of stock that are grazed on the near-by range.

The average annual rainfall in this section is about 11 inches, which usually is sufficient to produce crops on good soil which is well farmed. The principal drawback to crop production is the frequency of summer frosts. The nights are nearly always cool and frost may occur in any of the summer months. The frost-free period is seldom more than 40 or 50 days.

Winter wheat is the best of the cereal crops, especially on the valley silt loams and the plateau fine sandy loams. It should be sown on summer fallow at the rate of about 30 pounds to the acre. Only the hard red varieties, such as Turkey and Crimean, should be grown. When there is not sufficient moisture in the autumn to germinate winter wheat, spring wheat should be sown.

Rye is the principal crop now grown in central Oregon. Usually more grain is produced when it is sown in early spring than in the fall, for it is later in heading and more often escapes summer frosts.

Because of the distance from market and the low price per pound which can be obtained for oats and barley as compared with wheat, these grains should be grown only for feeding on the farms where they are produced. Sixty-Day oats and White Smyrna barley are the best varieties of these grains to grow.

GRAINS FOR THE DRY LANDS OF CENTRAL OREGON.

CONTENTS.

	Page.		Page.
Place of the grain crops in central Oregon	3	Small-grain crops	17
Description of central Oregon	4	Wheat	17
Equipment needed for grain growing on dry lands	9	Emmer and spelt	18
Rotation of crops	10	Rye	18
Preparing the land for grain	12	Oats	20
Good seed	13	Barley	20
Treatment of seed for smut	14	Flax	21
Pests	15	Rank of small-grain crops	21

PLACE OF THE GRAIN CROPS IN CENTRAL OREGON.

The future of the dry lands of central Oregon lies in a combination of crops with stock. The cultivated land will be used to produce feed for wintering and finishing stock that are grazed on the near-by range, only the surplus crops being sold. This will make dry farming succeed where otherwise it would be impossible.

In such a system of farming the growing of grains has a very important place. This bulletin therefore gives general information, based partly upon experiments at the Harney Branch Experiment Station at Burns, Oreg.,[1] concerning the production of the small grains on those nonirrigated lands of central Oregon which lie at altitudes between 4,000 and 5,000 feet. To lands which have the water table at a depth of 10 feet or less or which can be irrigated, the contents may apply in part. They apply only where the rainfall, summer frost, and winter cold are average and where the nature, depth, and alkali content of the soil are favorable for crop production. Market limitations and the needs of individual farmers are important factors also in determining the choice of crops among those that can be grown.

The important cereal crops for these dry-farming lands as a whole are winter and spring wheat, winter and spring rye, spring oats,

[1] The Harney Branch Experiment Station was established at Burns, Oreg., in 1911. Harney County appropriated funds for the purchase of land, the erection of buildings, and the purchase of equipment. The work is conducted cooperatively by the Oregon Agricultural Experiment Station and the United States Department of Agriculture. Three years' results have now been obtained on summer-fallowed land.

and spring barley. Flax, field peas, and dry-land alfalfa are also of importance, but their place is not yet fully determined. Crops of minor importance which can be grown are emmer, spelt, rape, sweet clover, root crops, and potatoes. Crops which have been rather definitely proved to be unadapted to the average conditions are corn, millet, sorghum, beans, buckwheat, and cultivated grasses. Any crop which requires more than a very short growing season and is not resistant to frost and drought should not be attempted in more than an experimental way.



FIG. 1.—Map of Oregon, showing the principal divisions of the State, with contour lines up to 6,000 feet east of the Cascade Range. The shaded portion of the map shows the section to which this bulletin particularly applies, that part of central Oregon with altitudes between 4,000 and 5,000 feet.

DESCRIPTION OF CENTRAL OREGON.

The section commonly known as central Oregon covers about one-third of the State. It includes the southern half of that part of Oregon lying east of the Cascade Range (fig. 1). Its total area is about 26,000,000 acres, of which possibly 5,000,000 acres are arable. Probably 1,000,000 acres of the arable land can be irrigated eventually in some way. The area that must be dry farmed, if farmed at all, is therefore about 4,000,000 acres. The land that is fit only for range amounts to about 21,000,000 acres. At present, railroad transportation has barely touched the edge of this great area and comparatively few people are making their homes upon the nonirrigated lands.

TOPOGRAPHY.

Central Oregon is a region of great plateaus broken by wide, level valleys and low, well-weathered mountains or buttes, with few high mountains and few streams. Most of the tillable dry-farming land lies between elevations of 4,000 and 5,000 feet. Central Oregon lies in that division of the United States known as the Great Basin, from which there is no drainage to the sea. Both the water and the cold air from the mountains flow down into the valleys.

SOILS.

The tillable dry-farming soils of central Oregon probably can be described best as the valley silt loams, valley sandy loams, valley sands, and plateau fine sandy loams.



FIG. 2.—A vigorous growth of black sage in the Harney Valley in central Oregon, indicating fertility without excess of alkali.

VALLEY SILT LOAMS.

The valley silt loams include the heavier valley soils. They occur in practically all the valleys, large or small, and occupy probably about one-third of the entire valley area. Usually they are gray, brown, or black in color. The subsoil usually is quite heavy for a few feet, giving way to a more open soil at a depth of 6 feet or more. Hardpan may or may not be present. Water is found at depths of 10 to 200 feet, according to location. The native growth varies with the alkali content and other soil variations. A uniform growth of black sage (Artemisia), such as is shown in figure 2, is a fair indication of fertility without excess of alkali. Salt-grass (Distichlis), greasewood (Sarcobatus), and rabbit brush (Bigelovia) should be looked upon as a warning that injurious alkali may be present.

VALLEY SANDY LOAMS.

Those soils which are light enough to be tilled easily but not so sandy that they are moved much by the wind are classed as valley sandy loams. They occupy probably half of the total valley area. They are yellowish or brownish in color. The subsoil usually is somewhat heavier than the surface soil for a few feet and then becomes lighter, like that of the valley silt loams. The depth to water and the native vegetation, alkali content, and fertility are similar to those of the valley silt loams.

VALLEY SANDS.

The valley sands include those soils which are so sandy in character that they are blown about by winds. They are very easily tilled but otherwise difficult to handle. They comprise probably about one-sixth of the total valley area. The native vegetation, depth to water, and alkali content are similar to those of the other valley soils.

PLATEAU FINE SANDY LOAMS.

The plateau fine sandy loams, occurring on the hills and plateaus, are easily tilled, brown in color, and vary greatly in depth and subsoil condition. Water usually is not found before a depth of 100 feet is reached, and sometimes not within 300 or 400 feet of the surface. The native growth is principally black sage or black sage and bunchgrass. These soils are fertile and seldom contain much alkali. The important thing, aside from the variation in climatic conditions with locality, is that they be deep and without hardpan. There should be 3 to 4 feet of good soil to make these sandy loams suitable for farming.

CLIMATE.

The successful production of crops in central Oregon depends more on the climate than on any other factor, even though the altitude is not great and the Pacific Ocean not distant.

RAINFALL.

The average annual rainfall over the whole of central Oregon is about 11 inches. Most of this comes during the winter and spring months. Very little rain falls during the growing season. The rainfall varies markedly, however, even within short distances. Ordinarily, most of the moisture is absorbed by the soil, but there are times in the spring when the loss by run-off is great. This is true particularly on the valley silt loams and the steeper slopes of the plateau fine sandy loams. Sufficient moisture usually falls to produce

crops on good soil, well farmed, in a locality having rainfall that is average or better. Snow lies on the ground for three or four months during the winter in some localities, thus affording excellent protection for hardy fall-sown grains. In other localities the ground remains covered for only a short time.

EVAPORATION.

The average evaporation from a free water surface at the Harney Branch Experiment Station at Burns is about 45 inches for the months from April to October, inclusive. Wind and heat increase the loss by evaporation. Good tillage and early-maturing crops are measures against crop failure. Evaporation during the winter months is slight.

WIND.

The average velocity of the wind is only about 4 miles an hour. At the Harney Branch Experiment Station it is highest in the spring months. The highest average velocity recorded at Burns for a 24-hour period in 1915 was 14.4 miles an hour on April 30. High winds, though not frequent, do considerable damage on the valley sandy soils by destroying young crops, cutting them off, blowing them out, or burying them when the sand moves with the wind.

TEMPERATURE.

The average daily mean temperature is about 44° F. This average daily mean temperature is from 5 to 10 degrees lower than that of the Columbia Basin, about 3 degrees lower than that of the Palouse section of eastern Washington and western Idaho, and from 2 degrees to 4 degrees lower than that of the Snake River basin of southern Idaho. It is about the same as the average daily mean temperature of eastern South Dakota and southern Minnesota. The average daily minimum temperature is about 27°, and the average daily maximum about 60° F. The highest summer temperature is rarely as much as 100° F., and the lowest winter temperature is rarely lower than —25° F. The days in summer are warm, but pleasant; with few exceptions the nights are cool.

The greatest problem in farming in central Oregon lies in the frequency of summer frosts. Except in favored places, summer frost rarely, if ever, fails to occur in June. It sometimes occurs in July, and seldom fails to occur before August 20. The favored places referred to are creek canyons and the slopes of hills where good air drainage is afforded or where the temperature is moderated by nearness to bodies of water. The average length of the growing season for such crops as corn is 40 to 50 days except in the comparatively small favored localities. Some crops are much hardier and under good management usually can be grown without serious damage.

AGRICULTURAL HISTORY.

The early settlers in central Oregon were stockmen. They selected land upon which wild grasses could be grown for hay to winter the stock that grazed on the ranges during the rest of the year. The ranges were overstocked and the native grasses were grazed so closely that they were largely destroyed. The annuals could not mature sufficient seed to reseed themselves, and the perennials were trampled out or pulled up. Reductions in the number of stock on the range came as the natural consequence.

The overgrazing of the ranges was followed by an influx of new settlers, who began fencing much of the range lands for dry farms. These new settlers did not engage in stock growing, but put their dependence on grain production. They were poorly equipped for

FIG. 3.—Mountain and hill land in Malheur County, Oreg. For every acre of tillable land in central Oregon there are many acres of this land that will always be range.

farming. The cost of living and of crop production far from railroads was high, climatic factors were adverse, and rodent pests were numerous. Many settlers attempted to produce grain without putting the land in shape to make production possible. They failed, but the range had opportunity to regain much that it had lost.

AGRICULTURAL POSSIBILITIES.

To-day the once boundless ranges of central Oregon are broken by the fences of settlers and by the remains of fences whose builders have gone. The day of the stockman who ranged his stock on a thousand hills and valleys is passing. With it is passing also the day of the man who depended on grain farming alone. It is a region of much Government land awaiting settlement, of much cheap

deeded land, and of great areas of reviving range land. The day of the small stockman-farmer is at hand.

Transportation has not yet and may not soon come within practical hauling distance by wagon of most of this land. Even when it does, it is doubtful whether grain farming alone will be a profitable form of agriculture over most of this section. For every acre of land that can be cultivated there are many acres of range land (fig. 3). Any system that does not utilize both the tillable land and the range can hardly be fully successful.

Dry farming, to be successful, must be done right, and the ranges must be brought under control so that they can not be destroyed by overgrazing. With adapted crops of good varieties, with correct methods of production in use on the tillable lands, and with some system of controlled or rotation grazing for the range, central Oregon will produce many times as much wealth in live stock as it has in the past.

EQUIPMENT NEEDED FOR GRAIN GROWING ON DRY LANDS.

A farmer may buy more machinery than is needed or he may get the wrong kind or not buy enough. The average farmer should depend upon horses rather than engines for power. Horses can be kept quite cheaply if grazed on the range, and the value of colts may make them a real asset rather than an expense. The delay and expense of getting repairs and the high cost of fuel, due to the necessity of a wagon haul of 100 to 200 miles from the railroad, make traction engines a questionable investment at the present time for the average dry farmer in central Oregon.

Every farmer should have a plow, a disk harrow, a spike-tooth harrow, and a drill. The disk plow often is better than one of the moldboard type for central Oregon land, because it works better through the sticky spots. Disk plows also are better for plowing dry land. However, the type of plow which is most desirable depends to a considerable extent on the locality. The machinery in successful use in a neighborhood is a good guide in buying. The double-action disk harrows are best, the cutaway type being good where the land crusts badly. Any well-made spike-tooth harrow is good. A spring-tooth harrow mounted on high wheels and equipped with interchangeable teeth and teeth points can perhaps be substituted for both the disk and spike-tooth harrows on the average farm. The single-disk drill is better for trashy land and the double-disk and hoe drills for clean land.

Often several farmers can combine in buying the more expensive machinery, thus lessening the individual expense. Sometimes, also, it is cheaper to hire machinery from another farmer or to pay him to do certain work than to buy expensive machinery which will seldom be used.

ROTATION OF CROPS.

In planning a crop rotation for dry-farmed land in central Oregon, two questions should be kept in mind: (1) What combination of crops will give the greatest immediate return? (2) What will be the effect upon soil moisture, fertility, and future crops? If only the immediate return is considered, those crops should be chosen which are best adapted to the individual farm, and they should be sown in the order that will give the greatest return at the least cost. While it is important to have the permanent productiveness of the soil in mind, immediate profit is far more important to the average dry farmer in central Oregon under present conditions.

The desire for present profits, however, should not cause one to lose sight of future possibilities. To maintain the fertility of the soil, the rotation should include nitrogen-gathering crops, like peas, alfalfa, and sweet clover. If the immediate profit is about equal to that from a rotation including cereals only, no one can afford to omit the legumes.

In the selection of a rotation the individual farmer must consider: (1) The product which will pay best, whether hay to feed range stock in winter, grain to sell, or both; (2) the crops most likely to give the greatest quantity of the products wanted; (3) the order in which these crops shall follow each other; and (4) the tillage which shall be given the land.

No rotation practice can succeed on these lands if it does not include clean summer tillage. Whether this summer tillage must be given every other year or only once in three or more years has not yet been demonstrated fully. However, it is known that on the average the second grain crop after the summer fallow will be light. Usually it will be a failure. A cultivated crop or a very early maturing cereal which may be cut for hay if it fails to make grain might do. More information must be obtained before any cultivated crop can be recommended generally. Only those rotations are discussed, therefore, which include summer fallowing once in two or three years.

As already stated, the grain-producing crops adapted to central Oregon in general are wheat, rye, oats, and barley, with peas as a possible leguminous seed crop. The crops important for hay are rye, wheat, and oats, in the order named, with field peas and sweet clover as possible leguminous hay crops suited to short rotations. Alfalfa can not be considered in short rotations, for the reason that it should be kept on the land several years when a stand is once established.

TWO-YEAR ROTATIONS.

The 2-year rotation, fallowing half the land each year and cropping the other half, is the safest for the farmer who grows only one crop, whether wheat, barley, or oats. On a 320-acre farm the arrangement each year would be simply 160 acres of summer fallow and 160 acres of one or more of these crops. Where two or more crops are grown they should change places on the land the next time the land is cropped. In this way the same kind of crop might not be grown on the same field more than once in four or even six or eight years. Whether or not the 2-year is better than a 3-year rotation containing one year of fallow, one grain crop, and one hay or cultivated crop depends on many local conditions. It can not be answered fully here.

THREE-YEAR ROTATIONS.

Under the 3-year rotation plan one-third of the land would be summer-fallowed each year, one-third cropped to grain following summer fallow, and one-third used for some crop sown on stubble land, double-disked in the spring. This plan has the advantage of requiring plowing only once in three years and also gives a greater acreage of crop each year. It is of doubtful value, however, because few crops will make a profitable yield following a grain crop in average years. A crop which will produce a fair yield regularly on the stubble land must be had, or the yields from 320 acres in a 3-year rotation may be less than from a 2-year rotation. If the total production is the same from each rotation the labor cost also will be about equal. The best plan, then, is the one which gives the greatest average total product from the land.

The crops suited to follow the grain crop in the 3-year rotation must either mature early or be capable of intertillage, or both. Those which may be used in this way are rye, field peas, and possibly sweet clover. Spring rye or winter rye sown very early in spring is preferable to fall-sown rye on stubble land. Peas and sweet clover, if it is possible to grow them profitably, would give a leguminous crop, and the peas a cultivated crop as well.

For the grain farmer, the best 3-year rotation apparently is one-third fallow, one-third wheat, rye, oats, or barley sown on summer fallow, and one-third spring-sown rye on disked grain stubble. If peas for seed can be grown, a portion or all of the well-disked or fall-plowed stubble land can be sown to this crop, thus changing the kind of crop from time to time. For farmers who want hay only, the choice of crops would be the same, except that sweet clover might possibly be used. The farmer desiring both hay and grain may use any combination of the suggested crops suited to his needs, his soil, and his climatic conditions.

XUM

PREPARING THE LAND FOR GRAIN.

CLEARING AND PLOWING.

The ordinary sagebrush land of central Oregon can be cleared during the summer months at a labor cost of about $2 per acre. This is done by the use of a drag drawn by six or eight horses. The brush usually is dragged both ways and then burned where it falls. If the brush can not be burned in this way, it is raked into windrows with heavy sagebrush rakes. Clearing by hand costs about $5 per acre for labor, except when the brush is dry enough to burn as it stands. The first plowing should be done as soon after clearing as possible, in order to prevent new growth and to make the period for the storing of moisture as long as possible. The labor cost of breaking land in central Oregon is $2.50 to $3 per acre.

When stubble land is to be plowed, the work usually should be done in the fall or early in the spring. To delay plowing until late spring or summer allows the weeds to grow, and the moisture needed for crops is lost. When plowing can not be done early, disking as soon as possible in the spring will assist in holding moisture. The depth of plowing should be varied slightly from year to year. The land need not be plowed oftener than once in two years if other tillage, such as disking, is substituted. On very light soils and with certain cropping systems discussed elsewhere in this bulletin, plowing may be done only once in three years when other efficient but less costly tillage operations are performed.

SUMMER FALLOW ON SUMMER TILLAGE.

The dry-farming soils in central Oregon can not be cropped to grain continuously. This has been well demonstrated through experiment and farm experience. If a crop is grown on the land every year, the return will not be sufficient to pay the cost of production except where there is seepage from higher slopes or in years of unusually heavy rainfall. It has been shown, however, that paying crops can be grown by cropping the land one year in two. The other year is devoted to accumulating moisture and fertility by means of the summer fallow.

At the Harney Branch Experiment Station spring wheat has produced an average yield of 6.5 bushels per acre by continuous cropping in a 4-year test. This was produced at a net loss of $1.65 per acre per year. In arriving at this figure it is considered that the wheat was worth 75 cents per bushel at the farm and that the cost of production was $6.55. The experiment showed that when the land was summer-fallowed every second year the average crop harvested was 16.5 bushels per acre. Using the same values but increasing the cost of production to $9.80 to cover summer tillage, the

higher harvesting and thrashing cost, and another year's interest and taxes, there remains a net gain of $2.57 for each acre cropped. On a 320-acre farm the returns under the first method would be $528 less than cost per year. Under the second method, where one-half the land is in crop each year, the returns are $411 per year above the cost of production, a difference of $939 per year in favor of the second method.

Summer tillage should prevent all weed growth and keep the soil surface rough and loose (fig. 4). This allows rain to be absorbed and prevents loss by run-off and evaporation. Stubble land should be plowed in the fall or early spring. Fall-plowed land and land that can not be plowed until late in the spring should be disked in early spring if possible, and then tilled with disk, spring-tooth, or

FIG. 4.—A perfect summer fallow, free from weeds, and with the surface soil rough and loose.

spike-tooth harrow only as necessary to prevent weed growth and keep the surface soil in good tilth. A loose, cloddy surface is ideal. A dusty mulch blows, prevents ready absorption of rain, and crusts. Wherever possible, plow at right angles to a slope. The seeding of fall crops usually can be done without extra preparation of the fallow. If the soil has become badly crusted it will be necessary to disk in the early spring before seeding spring grain.

GOOD SEED.

The value of good seed can not be overestimated. To be good, seed must be of good quality and free from weed seed and shriveled kernels. It must be of a variety which is adapted to the soil and cli-

matic conditions of the region, possessing earliness and hardiness. It must give the highest possible yield of the best quality. It must also be as free as possible from disease. A few additional cents per acre for good seed usually will be returned many times in the value of the crop. Specific statements as to varieties of the various grains which are adapted to this section and as to the best rates and dates of seeding are made in the discussion of the individual crops in the later pages of this bulletin.

TREATMENT OF SEED FOR SMUT.

Seed grain, unless definitely known to be free from smut, should be treated with a formaldehyde solution before seeding. The usual

FIG. 5.—Jack rabbits brought together in a drive near Burns, Oreg. Although more than 20,000 rabbits were killed at this place, apparently just as many as ever were present when the crops were ready for harvesting the following summer.

strength is 45 gallons of water to 1 pound of 40 per cent formaldehyde. The cleaned grain may be treated by sprinkling and shoveling until each kernel is moist. It may also be dipped and the smut balls skimmed from the surface of the liquid after stirring the grain to allow the smut to rise.[1] In either case the grain should be dried as quickly as possible. Care should be taken not to infect the seed again by using sacks or drills that have not been treated with formaldehyde. Always buy formaldehyde of standard strength, such as is supplied in sealed bottles.

[1] For a more complete discussion of the formaldehyde treatment for grain smut, see Farmers' Bulletin 507, "The Smuts of Wheat, Oats, Barley, and Corn," a copy of which may be obtained free on request to the Secretary of Agriculture, Washington, D. C.

PESTS.

Jack rabbits and ground squirrels, locally known as sage rats, are the most common animal pests in central Oregon. These pests are numerous enough to do great damage to growing crops in some localities. Many methods of killing jack rabbits have been tried, with different degrees of success. Poisoning is effective when carried on extensively over large areas during the winter months. Rabbit drives, trapping, shooting, and snaring are seldom effective, because they are not conducted on a sufficiently large scale to affect materially the total number of rabbits. Some idea of the number of rabbits in this section can be gained from figure 5, which shows a portion of those brought together in a drive at Burns. The bounty plan has been tried in two or more counties in central Oregon with



FIG. 6.—Grain field damaged by ground squirrels, locally known as sage rats. The early and liberal use of poison is the best insurance against loss of this kind.

partial success but at considerable cost. Whatever plan is adopted, it should be prosecuted vigorously and on an extensive scale. A single county is small enough for an effective unit. It would be much better if all the counties would combine on a plan. When the pests are reduced in numbers the efforts should be much increased rather than discontinued. A fight of this kind will cost a great deal, but the money will largely be returned in increased crops. Every interest in central Oregon would be benefited, for the eradication of this pest would mean much in the development of farming and the restoration of the range. Until the jack rabbits are eradicated the only safety for growing crops over much of the territory is in fencing with rabbit-proof wire. This wire fencing should be not more than 1¼

inches between line wires in the lower foot, 1½ inches in the next 6 inches, 2 inches in the next 6 inches, and 2½ or 3 inches in the upper foot of a fence 3 feet in height. A closely barbed wire should be sunk in the ground about 2 inches, to prevent digging under. The fence then should be placed over it, with the lower wire tight to the ground. Two barbed wires above the woven wire, one of which should be quite near it, will make the fence effective against stock. Such a fence will often pay for itself within a short time in crops saved, especially on the more isolated farms.

The work of sage rats in a grain field is shown in figure 6. These pests can best be kept in check by the use of poisoned grain. The following formula is recommended by the Bureau of Biological Survey:

Strychnin alkaloid (powdered)	1 ounce.
Baking soda	1 ounce.
Gloss starch paste (thin)	1 pint.
Corn sirup	One-fourth pint.
Glycerin	1 tablespoonful.
Saccharine	1 teaspoonful.
Oats	20 quarts.

Mix the powdered strychnin, soda, and paste to a smooth, creamy mass. Stir in the sirup, glycerin, and saccharine. Apply to the grain and mix until each kernel is coated; then dry the product. Keep all occupied burrows on the farm and adjoining lands supplied once every week by scattering about one teaspoonful of the poisoned grain about each opening. The use of larger quantities is wasteful and is also dangerous to live stock. Early poisoning is very important. A cheap effective poison liberally used will do much to keep this pest in check.

At the present time insect pests are not a serious problem in central Oregon. Those which occur occasionally in sufficient numbers to cause damage to crops are cutworms and grasshoppers.

Cutworms may be destroyed when they appear by sowing poisoned bran mash late in the evening along the borders of the area eaten down. The following formula for preparing poisoned bran mash is taken from Farmers' Bulletin 739 of the U. S. Department of Agriculture: Mix 50 pounds of wheat bran, 2 pounds of Paris green, and 6 finely chopped oranges or lemons. Then bring the whole mixture to the consistency of a stiff dough by the addition of low-grade molasses, such as is used for feeding to cattle, adding water when necessary. Distribute this bran over the infested field in small lumps. If bran can not be obtained readily, middlings or alfalfa meal may be substituted.

When grasshoppers become numerous, they may be destroyed by using a similar poisoned bait composed as follows: Wheat bran, 25

pounds; Paris green, 1 pound; white arsenic, 1 pound; lemons or oranges, 6 finely chopped fruits; low-grade molasses or "blackstrap," 2 quarts. Ordinary table sirup should not be used for this purpose, as it is not attractive to grasshoppers. In the dry-land regions, water should be added to this bait at the rate of 4 gallons to 25 pounds of the mixture, as the extra moisture is necessary in order to attract the grasshoppers. It is also important to distribute this bait in the late afternoon or early evening, just before the grasshoppers ascend the plants on which they usually pass the night.

SMALL-GRAIN CROPS.

WHEAT.

Wheat is the best yielding grain crop to grow on land alternately cropped and fallowed in central Oregon on soils adapted to its culture. The valley silt loams, the plateau fine sandy loams, and, to a less extent, the valley sandy loams, are most suitable.

WINTER WHEAT.

Under proper conditions winter wheat will outyield all other cereal crops. While worthy of trial elsewhere, it is particularly adapted to the valley silt loams and the plateau fine sandy loams, but only in localities where snow lies on the ground throughout the winter and where frosts do not occur too regularly in late June and in July. Yields should average about 20 bushels to the acre on land previously well summer-fallowed. Yields, however, will vary greatly from year to year, according to the damage done by frosts during the time of heading and filling.

The Turkey winter wheat is best adapted, because of its hardiness, good yields, and the high quality of the grain for milling purposes. Seeding should be done on summer fallow at the rate of about 30 pounds of good seed per acre. Drill rather shallow about September 10 if the soil is moist enough to sprout the seed. The wheat should make a good start in the fall if it is to survive the winter in good condition. If the soil is dry, it may be better to sow spring grain, though very late fall seeding will sometimes succeed. Harrowing in the spring after growth has started well may help the crop under some conditions and may hurt it under others. Soil types and seasons vary so much that this question must be settled by each farmer for himself.

SPRING WHEAT.

Spring wheat is probably the most consistent yielder of all the grain crops where soils and climatic conditions are suited to its production. The crop is adapted particularly to the valley silt loams,

the plateau fine sandy loams, and the heavier portions of the valley sandy loams. It is less likely to be injured by frost than winter wheat, because the period of heading and ripening comes later in the summer and does not last as long when early varieties are used. Sown on good summer fallow, it may be expected to average 15 to 20 bushels per acre. As a hay crop it hardly equals rye in yield under average conditions. The best variety now known for central Oregon is the Early Baart, which combines earliness with drought resistance and a high average yield of excellent milling wheat. Still better varieties may be found, but at present no mistake can be made in growing the Early Baart. For hay the Palouse Bluestem may be preferable, because it is beardless.

A spring-wheat variety must mature early to be adapted to extensive cultivation in central Oregon. Late varieties are frosted repeatedly before maturing; this results in low yields of shriveled wheat which produces flour of inferior quality. Good, plump wheat is necessary in order to assure a market.

It is poor policy for a community to grow a large number of varieties. If all farmers grow the same variety, the product will be more uniform, with consequent benefits in price.

Spring wheat should not be drilled deeper than is necessary to place the seed well into moist soil. Sow about April 15 to 20 at the rate of about 30 pounds of good seed per acre. It is not advisable to harrow the crop after it is up.

EMMER AND SPELT.

Winter emmer and spelt are not adapted to central Oregon conditions, because neither is hardy enough to survive the average winter. Spring emmer can be grown and will make fair yields, but it is not as good either in yield or in value per pound as wheat, oats, or barley. Until emmer and spelt have been tested more thoroughly, they should not be grown in this region unless in an experimental way.

RYE.

Rye is the principal crop now grown on the dry-farming lands of central Oregon. Probably it will continue to hold first rank in acreage for a long time, though wheat may yet outrank it in total production of grain.

WINTER RYE.

Winter rye is the only type of rye now grown in central Oregon to any considerable extent. The crop is sown both in the fall and in the early spring. When sown in the fall under proper conditions it usually survives the winter very well and makes an early growth in the spring. This earliness, however, unless checked by pasturing,

very frequently results in the loss of part or all of the grain yield because of the frosting of the heads. When the crop is to be harvested for hay, this fact is slightly less important. When winter rye is spring sown, the development is slower and good yields of both grain and hay are more uniformly obtained than from the fall-sown winter rye. However, if sown in the spring, it must be sown very early or it will not succeed. If sown at the usual time of sowing spring grains, it will rarely get beyond the stooling stage except in unusually wet and cool summers.

As a fall-sown crop, winter rye is adapted particularly to the valley silt loams and plateau fine sandy loams where June and July frosts are rare and where there is usually a winter covering of snow. Under these conditions on well-fallowed land it may be expected to yield 12 or 15 bushels of grain or 1½ to 2½ tons of hay per acre. Where wheat is the first crop sown after fallow, rye is better than other cereals to sow on the disked stubble land if such a practice is adopted. Seeding should be done about September 10 to 15 at the rate of 30 to 60 pounds of good seed per acre.

Winter rye sown in early spring will succeed under nearly all soil and climatic conditions in that part of central Oregon under discussion where any grain crop would succeed. However, it is not always possible to seed it early enough and a delay of one or two weeks often results in failure. As good average yields may be expected as from fall-sown winter rye under like soil conditions. Since a winter covering of snow is not essential with spring-sown winter rye, it may be grown over a much wider area. It should be shallow drilled not later than April 1 at the rate of 30 to 60 pounds of good seed per acre.

The value of rye grain for feeding purposes is not quite equal to that of wheat, barley, or oats, pound for pound. It usually sells on the market, however, for about as much per pound as wheat. As a hay crop for wintering stock from the range, the value of the hay will be in proportion to the number of stock that the grower can winter on the product of his farm. Under good management the hay might bring the grower as much as $7 to $8 per ton when fed on the farm.

SPRING RYE.

Little or no true spring rye is being grown in central Oregon. True spring rye develops very rapidly when sown in the spring. It does not stool as much as winter rye and makes a shorter and finer growth of straw. The grain is also somewhat smaller than that of winter rye.

Spring rye is being tested at the Harney Branch Experiment Station, but the trials have not been conducted long enough to warrant a positive statement as to the usefulness of the crop in that locality.

So far as soils and climate are concerned it is adapted to as much of the dry-farming area of central Oregon as any grain crop. Its value will probably be greatest on the lighter soils and as a grain or hay crop on disked wheat stubble when the wheat follows fallow. Yields of hay and grain may be expected to be as good as or slightly better than those of early spring-sown winter rye. The crop should be drilled about April 20 at the rate of 30 to 60 pounds of good seed per acre, placed as shallow as possible to get the seed well into moist soil.

OATS.

The cool climate of central Oregon is favorable to oats, but the crop is not so drought resistant there as wheat. As a producer of grain in pounds per acre, oats may be expected to rank second to wheat under average conditions upon fallowed land. Where oats are needed at home, there is reason for growing the crop. As a market crop or as a feed crop for fattening meat animals, it does not have as high a value per pound as wheat.

WINTER OATS.

Winter oats will not survive the ordinary winter in central Oregon. Sown in the spring it is doubtful whether the yield would be as good on the dry lands as that from a good spring variety.

SPRING OATS.

Spring oats are best adapted to the valley silt loams. In seasons of favorable summer precipitation they will do well on the valley sandy loams and plateau fine sandy loams. When an early variety is seeded after April 15 heading does not occur till July and the crop is mature before August 10, so that little damage from frost is likely. On the soils adapted to the crop an average yield of about 30 bushels of grain per acre may be expected on summer-fallowed land. As a hay crop an average of 1 to 1½ tons per acre would probably be harvested.

The early small-kerneled Sixty-Day variety of oats is recommended for dry-farmed lands in central Oregon, especially for the production of grain. Seeding should be done from April 15 to 20 at the rate of 1 bushel of well-cleaned, plump seed per acre, drilled as shallow as possible, but placed well into moist soil.

BARLEY.

Barley is probably less adapted to central Oregon dry-farming lands than wheat, oats, or rye. Barley will do very well where the soil is usually quite moist and in seasons of abnormally heavy summer rainfall. Present information indicates that on the average it is

not nearly as good as wheat for the production of grain and is slightly less valuable than oats.

WINTER BARLEY.

Winter barley will not survive the average winter in central Oregon. When it survives, June and July frosts usually catch the crop in head. When spring sown it has not given as good results as good spring varieties.

SPRING BARLEY.

The importance of spring barley lies principally in supplying special local demands. Sold on the market, it usually brings less per pound than wheat or oats. Fed to fattening meat animals, it is about equal to wheat in value per pound. The crop is adapted to the valley silt loams particularly and can be grown on lighter soils when the summer rainfall is favorable. Barley is subject to frost injury, requiring late seeding to prevent heading before the last spring frosts. An average grain yield of 18 to 20 bushels per acre may be expected on summer-fallowed land where the soil is adapted to the crop.

The White Smyrna variety is probably the best adapted to the average central Oregon dry farm. This variety is very early and of good quality. Its short straw is an advantage from the standpoint of grain production. Seeding should be done about May 1 at the rate of 40 pounds of good seed, drilled as shallow as possible, but deep enough to place the seed well into moist soil.

FLAX.

The growing of the flax crop is yet in the experimental stage in central Oregon. Experiments conducted during the last three years indicate that it is quite frost hardy and that it is adapted to the heavier soil types. With improved transportation facilities, flax is a possible cash crop on soils adapted to its growth for use in rotation with other crops. A 3-year average yield of 6.6 bushels per acre has been obtained at the Harney Branch Experiment Station. The price per bushel on the farm may be expected to average from $1.25 to $1.50.

The best variety of flax to grow in this section has not yet been determined. Seeding probably is best done early in May at the rate of 16 pounds per acre, drilled shallow in a fine, firm, moist seed bed.

RANK OF SMALL-GRAIN CROPS.

Any ranking of crops adapted to central Oregon according to their importance must be qualified with the understanding that the ranking applies to the region as a whole. Under local variations of

soil, rainfall, frost, winter snow covering, transportation facilities, range conditions, actual products wanted, etc., the ranking given may or may not hold.

From the standpoint of grain production the adapted cereal crops rank in importance in the following order: Wheat, rye, oats, barley.

From the standpoint of yield and value when cut for hay the adapted cereal crops rank as follows: Rye, wheat, oats, barley.

PUBLICATIONS OF THE U. S. DEPARTMENT OF AGRICULTURE TREATING OF CEREAL PRODUCTION AND DRY FARMING IN THE PACIFIC NORTHWEST.

AVAILABLE FOR FREE DISTRIBUTION.

Range Improvement by Deferred and Rotation Grazing. (Department Bulletin 34.)

Experiments with Spring Cereals at the Eastern Oregon Dry-Farming Substation, Moro, Oreg. (Department Bulletin 498.)

Oats: Growing the Crop. (Farmers' Bulletin 424.)

Barley: Growing the Crop. (Farmers' Bulletin 443.)

Sweet Clover. (Farmers' Bulletin 485.)

The Smuts of Wheat, Oats, Barley, and Corn. (Farmers' Bulletin 507.)

The Field Pea as a Forage Crop. (Farmers' Bulletin 690.)

Marquis Wheat. (Farmers' Bulletin 732.)

FOR SALE BY THE SUPERINTENDENT OF DOCUMENTS, GOVERNMENT PRINTING OFFICE, WASHINGTON, D. C.

Farm Experience with the Tractor. (Department Bulletin 174.) Price, 5 cents.

Experiments with Marquis Wheat. (Department Bulletin 400.) Price, 10 cents.

Leguminous Crops for Green Manuring. (Farmers' Bulletin 278.) Price, 5 cents.

The Repair of Farm Equipment. (Farmers' Bulletin 347.) Price, 5 cents.

Sixty-Day and Kherson Oats. (Farmers' Bulletin 395.) Price, 5 cents.

Oats: Distribution and Uses. (Farmers' Bulletin 420.) Price, 5 cents.

The Control of Blowing Soils. (Farmers' Bulletin 421.) Price, 5 cents.

Pasture and Grain Crops for Hogs in the Pacific Northwest. (Farmers' Bulletin 599.) Price, 5 cents.

○

INDEX.

	Farmers' Bulletin No.	Page.
Abandoned farms, use of sweet clover, note	797	3
Abortion, cattle—		
bulletin by Adolph Eichhorn and George M. Potter	790	1–12
extent and nature of disease	790	3–4
immunity	790	6–7
spread, means and preventive measures	790	6–12
symptoms, diagnosis and treatment	790	4–12
Accounts, farm, use of diary, bulletin by E. H. Thomson	782	1–19
Achoreutes armatum, description, occurrence and control in mushrooms	789	8–9
Acid soil, liming for sweet clover growing, experiments	797	14–17
Agaricus campestric, description, occurrence and edible value	796	11–12, 13
Agriculture Department, cooperation with Federal Loan Board	792	12
Alabama, fur-bearing animals, laws 1916	783	4
Alaska, fur-bearing animals, laws, 1916	783	4
Alberta, fur animals, laws, 1916	783	21–22
Alfalfa, growth on alkali soil after sweet clover	797	14
Alkali, resistance of sweet clover	797	13–14
Amanitas, mushrooms, species, descriptions and poisonous nature	796	4–5
Amortization table, various rates of interest	792	6
Anthracnose, cotton, symptoms, cause, and control	787	39
Anthrax—		
acutus, symptoms	784	5
bacilli, vitality, conditions favorable, etc	784	7–9
bulletin by Henry J. Washburn	784	1–16
description, cause, symptoms and control	784	1–16
diagnosis, distinction from blackleg and Texas fever	784	9–11
forms and symptoms	784	4–6
history, notes	784	3
peraculus, symptoms	784	4–5
subacute form, symptoms, appearance of carcass	784	5–6
Ants, control by carbon disulphid	799	14–15
Aphids—		
plant, control by carbon disulphid	799	16–18
underground, control by carbon disulphid	799	16–17
Aphiochaeta albidihalteris, description, occurrence on mushrooms and remedies	789	4–6
Argentine ants, control by carbon disulphid	799	14–15
Arizona, fur animals, laws, 1916	783	4–5
Arkansas, fur animals, laws, 1916	783	5
Armadillidium vulgare, description, occurrence on mushrooms and control	789	10
Armillaria Mellea, description	796	6
Army worm, control, furrow method	793	27
ASHBROOK, FRANK G., bulletin on "Castration of young pigs"	780	1–6
Bacillus—		
abortus, cause of contagious abortion in cows	790	3
anthrax, description	784	3–4, 8, 9
Bacterium malvacearum, cotton disease, symptoms and remedial measures	787	33–34

	Farmers' Bulletin No.	Page.
Bait, poison—		
grain pests, formulas	800	16–17
sowbugs in mushroom beds	789	11
Bajra, note	793	3
Banks, Federal Land—		
aid to credit of farmer	792	10–11
relation of Government	792	11–12
Barley—		
growing on dry lands of Oregon	800	20–21
value by acre, comparison with flax, wheat, and oats	785	6
varieties for Maryland and Virginia	786	19–20
Barnyard millet, note	793	4
BATES, CARLOS G., bulletin on "The windbreak as a farm asset"	788	1–16
Bean soup, feeding to calves, preparation and value	777	10
Bear, bounty paid	783	5, 7, 13, 19, 20
Beefsteak mushroom, description, edible value, and other names	796	16
Beetles, Lyctus powder-post, damage to seasoned hardwood	778	8–12
Belgian hare, wild, bounty	783	6
Bengal grass, description and characteristics	793	8–10
Billion-dollar grass, note	793	4
"Bird" cherry, grafting stock, note	776	6
Black turkey, description, weight, and value	791	5, 6
Blackbird, bounty	783	11
Blackhead, turkeys, symptoms and treatment	791	24–25
Blackleg, diagnosis, distinction from anthrax	784	9
Black-root, cotton, symptoms, cause, and control	787	35–38
Blue cotton, nature and remedies	787	40
Bobcat, bounty	783	9, 19, 20
Boletus mushroom, description and doubtful value	796	15
Boletus, spp. occurrence and description	796	15
Bonds, issue by Federal land banks	792	7
Bookkeeping, farm accounts, use of diary	782	1–19
Borers, wood, control by carbon disulphid	799	19
Bounties, wild animals, laws by States and Provinces	783	4–25
BRAINERD, W. K., and H. P. DAVIS, bulletin on "Feeding and management of dairy calves and young dairy stock"	777	1–20
Breeding, turkeys, selection of variety and stock, management, etc.	791	6–13
BREITHAUPT, L. R., bulletin on "Grains for the dry lands of central Oregon"	800	1–22
British Columbia, fur animals, laws, 1916	783	22
British West Indies, production of Sea Island cotton	787	8
Bronze turkey, description, weight, and value	791	5, 6
Brooding, turkey poults	791	16–21
Broom-corn millet, notes	793	3, 4
Bud variation, citrus trees, occurrence and control	794	4
Bud wood, citrus trees, selection and care	794	12, 14
Budding, citrus—		
selection of buds to avoid bud variations	794	4
trees, directions	794	14
Buhach, use against springtails on mushrooms	789	9
Buildings, fumigation with carbon disulphid, directions	799	12–14
Bull—		
breeding age, care, and handling	777	19
transmission of abortion diseases, treatment, etc	790	11
Buttermilk, creamery, carrier of tuberculosis to hogs	781	7
Calf—		
diseases, nature, and treatment	777	16–17
pens, management, bedding, and care	777	12
teaching to drink	777	3–4
California, fur animals, laws, 1916	783	5

	Farmers' Bulletin No.	Page.
Calvatis gigantea, note	796	17, 19
Calves—		
dairy, feeding, and management, bulletin by W. K. Brainerd and H. P. Davis	777	1–20
feeding at various ages, recommendations	777	17–18
horns, treatment to prevent development	777	14
marking for identification, practices and suggestions	777	14–16
Canada, fur-bearing animals, laws, 1916	783	2–3, 21–25
Canning mushrooms, methods	796	23–24
Cantharellus cibarius, description, occurrence, and edible value	796	7–8
Caponizing, turkeys, advantages	791	22
Carbolic acid, use against abortion of cows	790	12
Carbon—		
bisulphid—		
handling and storing, precautions	799	6–7
insecticide, bulletin by W. E. Hinds	799	1–21
properties, liquid and vapor	799	4–15
vapor, properties, effects, diffusion, etc	799	5–6, 8–10
disulphid—		
use against mushroom pests	789	5
variant names	799	2
Carcasses, anthrax infected, disposal	784	16
Castration, pigs, bulletin by Frank G. Ashbrook	780	1–6
Catch crop—		
millet, value	793	6
Sudan grass, value	793	6
Cats, anthrax, symptoms	784	6
Cat-tail millet, note	793	4
Cattle—		
contagious abortion, bulletin by Adolph Eichorn and George M. Potter	790	1–12
feeding millet hay	793	20
herd management against abortion disease	790	8–10
susceptibility to anthrax, dangers of swampy ground	784	6, 7–9
tuberculous, source of disease in hogs	781	3
Cedars, use in windbreaks	788	4, 13
Chaetochloa italica. *See* Millet, foxtail.		
Chanterelle mushroom, description and edible value	796	7–8
Charbon—		
bulletin by Henry J. Washburn	784	1–16
See also Anthrax.		
CHARLES, VERA K., and FLORA W. PATTERSON, bulletin on "Some common edible and poisonous mushrooms"	796	1–24
Cherries—		
diseases and pests, and control measures	776	23–25
growing east of Rocky Mountains, bulletin by H. P. Gould	776	1–37
picking and handling	776	25–29
protection from birds	776	24–25
ripening, sequence of different varieties	776	35–36
sour—		
growing, distribution and extent	776	4
varieties, descriptions and value	776	30–31
sweet—		
growing, extent and distribution	776	4
sterility, note	776	35
varieties, descriptions and value	776	32–34
varieties, descriptions and value	776	29–34
Cherry—		
growing, extent and distribution	776	3–4
industry, outlook	776	36
orchard, interplanted crops, suggestions	776	15–16
orchards, location and requirements	776	4–5

	Farmers' Bulletin No.	Page.
Cherry—Continued.		
trees—		
propagation and stocks	776	5–6
pruning	776	12–13, 16–23
selection, handling, planting, etc	776	6–13
Chicken pox, turkeys, symptoms and treatment	791	25
Chickens, anthrax, symptoms	784	8
Chinch bugs, control, furrow method	793	27
Citrus trees—		
records of crops, management	794	5–12
top-working	794	12–14
Citrus-fruit, improvement: How to secure and use tree-performance records, bulletin by A. D. Shamel	794	1–16
CLARK, CHARLES H., bulletin on "Seed-flax production"	785	1–20
Clavariaceæ, characteristics	796	16
Clearing, new land, central Oregon, cost and management	800	12–13
Clitocybe illudens, description and occurrence	796	6, 8
Clover, sweet—		
eradication	797	33–34
growing in mixtures	797	32–33
growing the crop, bulletin by H. S. Coe	797	1–34
history and distribution	797	10–12
inoculation, need and methods	797	27–29
seeding rate and methods	797	22–27
species, description	797	4–10
stand, treatment first and second seasons	797	30–32
white, description	797	5–9
yellow, biennial and annual	797	9
COE, H. S., bulletin on "Sweet clover: Growing the crop"	797	1–34
Colletotrichum gossypii, cause of anthracnose in cotton	787	39
Colorado—		
fur animals, laws, 1916	783	5
grass, note	793	4
Colostrum, value for calf	777	3–4
Compost, mushroom, sterilization	789	5
Connecticut, fur animals, laws, 1916	783	5–6
Contagious abortion of cattle, bulletin by Adolph Eichhorn and George M. Potter	790	1–12
Cooking mushrooms, recipes	796	19–23
Coprinus, species, descriptions, occurrence and edible value	796	12–14
Coracan, note	793	4
Coral fungi, description and warning	796	16
Corn—		
fumigation, with carbon disulphid, for insect control	799	11
growing, Minnesota, protection by windbreak, instance	788	4–5
husking in field for control of insects	799	11
protection by windbreak, effect on yield	788	10–11
yield, effect of windbreaks	788	5, 10–11
Corrals, sheep, in dipping work	798	25
Cortinarius mushroom, description, edible and poisonous qualities	796	11
Cotton—		
defects influencing grade	787	28–32
production, cost, effects of economic conditions	787	8–9
qualities desirable	787	25–28
Sea Island—		
bulletin by W. A. Orton	787	1–40
classification and preparation for market	787	5–6
cultivation, fertilizers, and rotations	787	9–18
distribution, crop requirements, and extension of area	787	2–5
history, characteristics and value	787	1–2
picking, ginning, and marketing, disadvantages	787	5
varieties, sale of seed, etc	787	19–20

	Farmers' Bulletin No.	Page.
Cottonseed—		
fumigation with carbon disulphid for insect control	799	14
Sea Island, selection	787	18–28
Cottonwood, use in windbreak planting, recommendations and returns	788	4, 11, 13, 14
Cougar, bounty	783	17, 22
Cow, abortion disease, treatment	790	12
Cowpeas, growing in mixtures for dairy cows, practices	793	21–23
Cows—		
contagious abortion, bulletin by Adolph Eichorn and George M. Potter	890	1–12
feeding—		
before parturition	777	3
mixture of millet and legumes, practices	793	21–23
pregnant, treatment	790	9
Coyote, bounty paid, by States	783	5, 7–8, 12–13, 16–22, 25
Creameries, tuberculous milk, per cent and distribution	781	6–7
Credit, farmer—		
benefits of Federal land bank	792	10–11
benefits of local farm-loan association	792	9–10
Creosote, coal-tar, dips for sheep tick	789	19
Cresol, dips for sheep ticks	798	19–20
Cresolis compositus, use against abortion of cows	790	11–12
Cresylic acid, per cent in sheep dips	798	18, 19
Cribbers, symptoms and objections	779	3
Crickets—		
injury to mushrooms, remedies	789	12
mole, control by carbon disulphid	799	15
Crop impaction, turkeys, cause and treatment	791	26
Crops, protection by windbreaks, yields, etc	788	3–8, 9–11
Crow, bounty	783	7, 8, 11, 20
Cutworms, poisoning	800	16
Dairy—		
calves, feeding and management, bulletin by W. K. Brainerd and H. P. Davis	777	1–20
cattle, young, feeding and management, bulletin by W. K. Brainerd and H. P. Davis	777	1–20
Dakota millet, description and characteristics	793	7–8
DAVIS, H. P., and W. K. BRAINERD, bulletin on "Feeding and management of dairy calves and young dairy stock"	777	1–20
Dead animals, anthrax infected, disposal	784	16
DEARBORN, NED, bulletin on "The Domesticated silver fox"	795	1–32
Death cup, mushroom, description and warning	796	5
Delaware—		
fur animals, laws, 1916	783	6
grain acreage, by varieties, 1911–1915	786	3
grain production, yields of various kinds, 1906–1915	786	21
Dairy, farm accounts, bulletin by E. H. Thomson	782	1–19
Diaries, types for farm accounts	782	4–6
Dipping—		
bag, sheep, use in tick control	798	23–24
forks, description and use	798	16–17
sheep—		
cost	798	17
directions	798	11–23
equipment and permanent plants, construction, etc	798	23–31
portable and permanent vats	798	26–29, 30–31
tick eradication, management and value	798	11–31
Dips, sheep, for ticks, formulas and mixing	798	18–23

	Farmers' Bulletin No.	Page.
Disinfectants, use against contagious abortion	790	10–11
Disinfection, tuberculous premises	781	16
Dogs, anthrax, symptoms	784	6
Dried mushrooms, preparation and uses	796	24
Dry lands—		
clearing and plowing, central Oregon	800	12–13
Oregon, grains suitable	800	1–22
summer tillage for grain in Oregon	800	12–13
Drying, mushrooms, methods	796	24
Duke cherries—		
descriptions and value	776	3
nature, growth, habits and pruning methods	776	21–22
Eagle, bounty	783	22
Early Baart wheat, characteristics and value for dry lands of Oregon	800	17–18
Early Harvest millet, description and characteristics	793	7–8
Echinochloa—		
colona, note	793	4
frumentacea, note	793	4
Eggs, turkey—		
care and incubation	791	13–16
production, care, and incubation	791	10–16
Egyptian—		
cotton, production and characteristics	787	7
millet, note	793	4
EICHHORN, ADOLPH, and GEORGE M. POTTER, bulletin on "Contagious abortion of cattle"	790	1–12
Eleusine coracana, note	793	4
Emmer, varieties for Maryland and Virginia	786	16–17
English sparrows, bounty	783	16, 21
Equestrian tricholoma, mushroom, description, edible value	796	9
European Lyctus, description, occurrence and distribution	778	12–13
Evergreens, planting for windbrakes, returns	788	12,13,14,15
Eyes, horse, examination for defects, suggestions	779	9–10
Fairy-ring mushroom, description and edible value	796	10
"Famine crop," use of millet in India	793	4
Farm—		
accounts, use of dairy, bulletin by E. H. Thomson	782	1–19
business, financial summary, records	782	7–8
Loan—		
act, Federal, benefit to farmer, bulletin by C. W. Thompson	792	1–12
association, National, organization of local body	792	8–9
Board, cooperation of Agriculture Department	792	12
Board, Federal, work	792	1
operations, diary, use and value	782	1–19
windbrake as asset, bulletin by Carlos G. Bates	788	1–16
Farmer, benefit from Federal farm loan act, bulletin by C. W. Thompson	792	1–12
Farm-mortgage loan system, disadvantages	792	4–5
Farms, abandoned, value of sweet clover, note	797	3
Feeding, foxes, directions and care	795	17–19
Fence, rabbit proof, use in central Oregon	800	15–16
Fences, fox yard, construction	795	11–19
Fertilizers—		
grains, fall-sown	786	5–7
millet, experiments	793	25–26
Sea Island cotton, selection and application	787	12–15
sweet clover, choice and application	797	17–18
Finger millet, note	793	4
Fistula, horse, symptoms, comparison with collar sores	779	12
Fistulina hepatica, description, occurrence and edible value	796	16

	Farmer's Bulletin No.	Page.
Flax—		
growing—		
on dry lands of Oregon	800	21
rotations	785	11–13
harvesting, thrashing and storage	785	16–19
types, descriptions and value	785	7
value by acre, comparison with wheat, oats and barley	785	6
Flax-growing area	785	5
Flaxseed—		
brushers, supply	785	4
center of production, changes	785	3
cleaning and treatment for wilt	785	9–10
consumption and production in United States, 1905–1914	785	3–4
industry, importance	785	4–5
markets	785	6–7
preparation for calves, nature and value	777	10–11
production, bulletin by Charles H. Clark	785	1–20
production for sowing, suggestions	785	8–10
production, 1905–1914	785	3
sowing, preparation of land, etc	785	13–16
Florida, fur animals, laws, 1916	783	6
Fly agaric, mushroom, description and warning	796	5
Food, foxes, selection and preparation	795	17–19
Forage, publications, list	797	35
Forks, dipping, description and use	798	16–17
Formaldehyde, use against grain smuts	800	14
Formalin—		
treatment of millet seed for smut	793	26
use against smut of millets	793	26
Fox—		
bounty paid	783	5, 13, 17–19
dens, types and construction	795	9–13
farming, history and growth of industry	795	5–7
food and feeding	795	17–19
killing for pelt, management	795	30–31
ranch—		
accessories	795	28–29
establishment, cost and profits	795	31–32
establishment, site and equipment	795	7–17
runs, fencing and plans	795	11–17
silver—		
domesticated, bulletin by Ned Dearborn	795	1–32
relation to red and black foxes	795	4–5
Foxes—		
diseases and parasites, treatment	795	22–25
domestication, history, requirements, and management	795	2–32
improved strains	795	25–28
marketing for breeding	795	29–31
propagation, habits, handling, etc	795	20–25
shipping, directions	795	23
Foxtail millet—		
culture and utilization, bulletin by H. N. Vinall	793	1–28
varieties, comparison	793	15–16
See also Millet.		
Frost, summer, central Oregon, relation to grain farming	800	2
Fumigation—		
bin, directions for making	799	8
buildings, with carbon disulphid, directions	799	12–14
cottonseed, with carbon disulphid, for insect control	799	14
mushroom houses, maggot control	789	4–6
Fur-bearing animals—		
laws relating to, 1916, bulletin by D. E. Lantz	783	1–28
trapping, open seasons, by States and Provinces	783	27–28
Fusarium vasinfectum, cause of cotton wilt	787	35–38

	Farmer's Bulletin No.	Page.
Garbage, feed for hogs, danger from tuberculosis	781	9–10
Georgia, fur animals, laws, 1916	783	6
German millet—		
description and characteristics	793	8–10
description and value	793	8–10
"Giant millet," note	793	4
Ginning, Sea Island cotton, disadvantages	787	5
Gold Mine millet, comparison with common millet	793	8
Golden millet, description and characteristics	793	8–10
Golden Wonder millet, description and characteristics	793	10
Gopher, bounty paid	783	5, 8, 11, 21
Gossypium barbadense. *See* Cotton, Sea Island.		
GOULD, H. P., bulletin on "Growing cherries east of the Rocky Mountains"	776	1–37
Grain—		
crop, millet, yields and uses	793	23–24
crops, protection by windbreaks	788	9–10
feeding to calves, preparation and mixtures	777	9–10
fumigation with carbon bisulphid	799	10–12, 21
growing, Oregon dry lands, equipment, crop rotations and preparation of land	800	9–13
harvesting, Maryland and Virginia, shocking and stacking	786	12–14
seed-bed preparation	786	8–9
varieties, selection for Maryland and Virginia	786	14
yields of various kinds in South Atlantic States	786	21–22
Grains—		
dry lands of central Oregon, bulletin by L. R. Breithaupt	800	1–22
fall-sown, in Maryland and Virginia, bulletin by T. R. Stanton	786	1–24
fertilizers, in Maryland and Virginia	786	5–7
Grape fruit, trees, records, and improvement in fruit	794	1–16
Grasshopper, control, formula	793	27–28
Grasshoppers—		
damage to millet, destructive measures, etc	793	27
poisoning	800	16–17
Green California millet, description and characteristics	793	7–8
Ground—		
hog, bounty	783	7, 11, 16
squirrel, bounty	783	11
squirrels, grain pest in Central Oregon, control methods	800	16
Grubs, white, control by carbon disulphid	799	15
Hardwood—		
products, injury by powder-post beetles, classes of articles	778	4
seasoned, damaged by Lyctus beetles, bulletin by A. D. Hopkins and T. E. Snyder	778	1–20
Hardwoods—		
seasoned, losses from powder-post beetle, extent	778	7–8
subject to powder-post damage	778	4
Harney Branch Experiment Station, Burns, Oreg., establishment	800	3
Harvesting—		
cherries, practices and suggestions	776	25–29
flax for seed	785	16–17
grain, Maryland and Virginia	786	12, 14
Hawaii, fur animals, laws, 1916	783	7
Hawk, bounty	783	7, 16–19, 21
Hay—		
anthrax transmission, note	784	9
millet—		
feeding value	793	20–21
harvesting, curing, and storage	793	19
injury to horses	793	21
value and yield of different varieties	793	8, 12, 13, 15–16

	Farmers' bulletin No.	Page.
Hayward's calf meal, ingredients and value	777	11
Hedgehog, bounty	783	13, 19
Heifers, breeding age, care, and handling	777	18–19
Heterodera radicicola, cause of root-knot of cotton	787	38
"High life," nature and use as insecticide	799	2–21
HINDS, W. E., bulletin on "Carbon disulphid as an insecticide"	799	1–21
Hog millet, note	793	4
Hogs—		
danger from tuberculosis in following cattle	781	7–10
prices, 1900–1909, 1912–1914	781	3
susceptibility to anthrax	784	6
tuberculosis—		
bulletin by John R. Mohler and Henry J. Washburn	781	1–20
infection sources	781	5–10
symptoms	781	10–11
See also Pigs.		
Honey-colored mushrooms, description and distribution	796	6
HOPKINS, A. D., and T. E. SNYDER, bulletin on "Powder-post damage by Lyctus beetles to seasoned hardwood"	778	1–20
Horns, development in calves, preventive treatment	777	14
Horse—		
foreparts, conformation and condition, requirements	779	7–18
gaits, and action, considerations	779	23–25
hind parts, conformation and condition, requirements	779	17–23
judging, considerations	779	5–25
parts, diagram	779	4
sound, selection, bulletin by H. H. Reese	779	1–27
Horses—		
feeding, millet hay, dangers and injurious results	793	21
injury from millet hay, experiments	793	21
susceptibility to anthrax	784	6
Household insects, control by carbon disulphid	799	19–20
Hughes, Prof. H. D., patentee of Ames scarifying machine for public use	797	20
Hungarian—		
grass, description and characteristics	793	10–12
millet, introduction and characteristics	793	10–12
Hunting, fur-bearing animals, laws, by States and Provinces	783	4–28
Hydrocyanic-acid gas, use against sowbugs in mushroom beds	789	11
Idaho, fur animals, laws, 1916	783	7
Illinois, fur animals, laws, 1916	783	7
IMES, MARION, bulletin on "The sheep tick and its eradication by dipping"	798	1–31
Incubation, turkeys	791	13–16
Indiana, fur animals, laws, 1916	783	7
Inoculation, sweet clover, need and methods	797	27–29
Insecticide, carbon disulphid, bulletin by W. E. Hinds	799	1–21
Insects—		
household, control by carbon disulphid	799	19–20
injurious, stored grain, control by carbon bisulphid	799	1, 10–12, 21
killing by carbon disulphid	799	1, 10–12, 21
Inventory, farm, suggestions for keeping	782	8–16
Iowa, fur animals, laws, 1916	783	7–8
Irrigation, flax crop	785	16
Italian millet—		
description and characteristics	793	7–8
note	793	4
Jack rabbit, bounty	783	5, 18
Jack rabbits, pest in central Oregon, damage to crops and methods of destruction	800	15–16

	Farmers' bulletin No.	Page.
Japanese millet—		
change of name to Aino millet	793	14
note	793	4
use of term	793	14
Jungle rice, note	793	4
Kansas, fur animals, laws, 1916	783	8
Ked. *See* Tick, sheep.		
Kentucky, fur animals, laws, 1916	783	8
Koda millet, note	793	4
Kursk millet, introduction and characteristics	793	13
Kutki millet, note	793	4
Labor—		
income, farmer, calculation	782	15–16
picking Sea Island cotton, prices	787	5
Sea Island cotton production, considerations	787	4–5
Lactarious mushrooms, descriptions and value	796	9–10
Lactarius, species, descriptions, occurrence, and edible value	796	9–10
Land, new, clearing and plowing, Oregon dry lands	800	12–13
Land-bank system, Federal, advantages	792	5–8
LANTZ, D. E., bulletin on "Laws relating to fur-bearing animals, 1916. A summary of laws in the United States and Canada relating to trapping, protection, propagation, and bounties"	783	1–28
Laws, fur-bearing animals, 1916, bulletin by D. E. Lantz	783	1–28
Legislation, fur-bearing animals, 1916	783	2–3
Lemon trees, records and improvement in fruit	794	1–16
Lepiota, species, descriptions and occurrence	796	6
Lice, extermination on turkeys	791	26
Limax maximus, description, occurrence on mushrooms and control	789	11–12
Limber neck, turkeys, symptoms, cause, and treatment	791	26
Lime—		
need in growing sweet clover, on acid soils	797	14–17
trees, records and improvement in fruit	794	1–16
use on millet, experiments	793	26
Lime-sulphur-arsenic dip, making, directions	798	20–22
Lindsey's calf meal, ingredients and value	777	11
Lion, mountain, bounty paid, by States	783	5, 7, 12–13, 17–21
Little millet, note	793	4
Loans—		
benefits of Federal farm loan act to the farmer, bulletin by C. W. Thompson	792	1–12
Federal land banks, requirements	792	8
Locust, use in windbreak planting, recommendations and returns	788	4, 6, 10, 12, 13, 14
Louisiana, fur animals, laws, 1916	783	8
Louse fly. *See* Tick, sheep.		
Lycoperdon pyriforme, description and value	796	17
Lyctus beetles—		
control measures	778	18–20
description, habits, and seasonal history	778	8–12
insect enemies	778	17–18
powder-post damage to seasoned hardwoods, bulletin by A. D. Hopkins and T. E. Snyder	778	1–20
species destructive to hardwood products, descriptions	778	12–15
Lynx, bounty, paid, by States and by Provinces	783	5, 7, 10–13, 17–21, 23
Mahaleb cherry, stock for grafting, notes	776	6
Maine, fur animals, laws, 1916	783	8–9
Man, anthrax, symptoms and means of infection	784	6–7

	Farmers' Bulletin No.	Page
Manitoba, fur animals, laws, 1916	783	22–23
Maple, planting in windbreak, returns	788	15
Marasmius oreades, description, occurrence, and edible value	796	10
Marketing—		
foxes for breeding	795	29–31
pelts of fox, history, preparation, and prices	795	5–7
Sea Island cotton, difficulties and suggestions	787	5–6
Markets—		
flaxseed	785	6–7
Sea Island cotton, location and grades	787	5–6
Marking calves for identification	777	14–16
Maryland—		
fall-sown grains, and in Virginia, bulletin by T. R. Stanton	786	1–24
fur animals, laws, 1916	783	9–10
grain—		
acreage, by varieties, 1911–1915	786	3
growing, fall, sowing, time, rate, and varieties	786	1–24
production, yields of various kinds, 1906–1915	786	21
Massachusetts, fur animals, laws, 1916	783	10
Mazzard cherry, stock, advantages for grafting	776	6
Melophagus ovinus. *See* Tick, sheep.		
Merulius lacrymans, description, occurrence, and injury to timber	796	16
Michigan, fur animals, laws, 1916	783	10–11
Milk—		
feeding to calves, quality and quantity	777	4–8
substitutes in feeding calves, preparation and value	777	10–12
tuberculous—		
danger to hogs	781	5–7
treatment for feeding hogs	781	7
Millet—		
diseases and insects injurious	793	26–28
feed value, and danger in feeding horses	793	20–21
food use in China	793	3
foxtail—		
culture and utilization, bulletin by H. N. Vinall	793	1–28
varieties	793	7–15
grain crop, yields and uses	793	23–24
growing in mixtures—		
for dairy cows	793	21–23
with legumes	793	21–23
history, climatic and soil requirements, etc	793	5–6
insect enemies, control	793	27–28
seeding, date, rate, and method	793	17–19
uses and value, yields, as grain crop	793	23–24
value in rotations	793	24–25
Millets—		
foxtail, description and characteristics	793	4–5
groups, common and botanical names	793	4
production and use in foreign countries	793	3–4
Minnesota, fur animals, laws, 1916	783	11
Mississippi, fur animals, laws, 1916	783	11
Missouri—		
fur animals, laws, 1916	783	11–12
millet, description and characteristics	793	7–8
Mite, mushroom, description, occurrence, and control	789	6–8
Moha de Hongrie, description and characteristics	793	10–12
MOHLER, JOHN R., and HENRY J. WASHBURN, bulletin on "Tuberculosis of hogs"	781	1–20
Mole crickets, control by carbon disulphid	799	15
Montana, fur animals, laws, 1916	783	12
Montmorency cherry—		
nature, ripening season, and value	776	30–31
trees, growth habits and pruning	776	18–19

	Farmers' Bulletin No.	Page.
Morchella esculenta, description, occurrence, and edible value	796	18, 20
Morel fungus, description and edible value	796	18
Morello cherry, nature, ripening season, and value	776	31
Moths, clothes, control by carbon disulphid	799	19–20
Mountain lion, bounty paid, by States	783	5,7,12–13, 18–21
Mulberry, planting in windbreaks, returns	788	14
Mules, susceptibility to anthrax	784	6
Museum pests, destruction by carbon disulphid	799	20
Mushroom—		
cultivated species, description, occurrence, and edible value	796	11–12, 13
fungi, varieties and descriptions	796	14–18
maggots, description, life history, and control	789	3–6, 11
mite, description, occurrence and control	789	6–8
pests and how to control them, bulletin by C. H. Popenoe	789	1–16
Mushrooms—		
cooking, receipes	796	19–23
edible and poisonous—		
bulletin by Flora W. Patterson and Vera K. Charles	796	1–24
detection methods, warning	796	2
edible, descriptions	796	6, 6–8, 9–14
food uses, poisoning, etc., precautionary measures	796	18–19
poisonous, descriptions and warnings	796	4–5, 8, 11
structure and descriptions	796	3–14
structure, and reproduction	796	3–4
Muskrat, bounty	783	20
Napoleon cherry, description and value	776	32–33
Narragansett turkey, description, weight and value	791	5, 6
Nebraska, fur animals, laws, 1916	783	12
Nests, turkey, preparation	791	12
Nevada, fur animals, laws, 1916	783	12
New Brunswick, fur animals, laws, 1916	783	23
New Hampshire, fur animals, laws, 1916	783	12–13
New Jersey, fur animals, laws, 1916	783	13
New land, flax growing, advantages	785	3, 11–12
New Mexico, fur animals, laws, 1916	783	13
New York, fur animals, laws, 1916	783	13–14
Newfoundland, fur animals, laws, 1916	783	23
Nicotin, dips for sheep ticks	798	20
Nicotine fumigation, use against mushroom pests	789	5
North Carolina, fur animals, laws, 1916	783	14–16
North Dakota, fur animals, laws, 1916	783	16
Northwest Territory, fur animals, laws, 1916	783	23
Nova Scotia, fur animals, laws, 1916	783	24
Nurse crop, sweet clover, need	797	18
Oats—		
fall sowing, time, method and rate, Maryland and Virginia	786	11, 12
growing on dry lands of Oregon	800	20
value, by acre, comparison with flax, wheat, and barley	785	6
varieties for Maryland and Virginia	786	18–19
Offal, slaughterhouse, feed for hogs, danger from tuberculosis	781	8
Ohio, fur animals, laws, 1916	783	16
Oklahoma, fur animals, laws, 1916	783	16
Ontario, fur animals, laws, 1916	783	24
Orange trees, records and improvement in fruit	794	1–16
Orchard, cherry—		
location and requirements	776	4–5
tillage, and soil maintenance	776	14–15
Orchards—		
citrus, treatment for citrus-fruit improvement	794	1–16
protection by windbreaks, advantages	788	8

	Farmers' Bulletin No.	Page.
Oregon—		
central, topography, soils, climate and agricultural possibilities.	800	4–9
fur animals, laws, 1916	783	16–17
grains for dry lands, bulletin by L. R. Breithaupt	800	1–22
ORTON, W. A., bulletin on "Sea Island cotton"	787	1–40
Osage orange, use in windbreak planting, recommendations and returns	788	12, 13, 14
Owl, bounty	783	7, 18–20
Oyster mushroom, description and edible value	796	7
Panicum—		
frumentaceum, note	793	4
germanicum, description and characteristics	793	10–12
miliaceum, note	793	4
miliare, note	793	4
psilopodium, note	793	4
texanum, note	793	4
Panther, bounty paid, by States	783	5, 14, 17–18
Parasol mushroom, description	796	6
Paris green, use against sowbugs in mushroom beds	789	11
Paspalum scorbiculatum, note	793	4
Pasteur vaccine, use against anthrax in sheep	784	12–13
Pasturing, millet—		
disadvantages	793	24
value for young stock	793	24
PATTERSON, FLORA W., and VERA K. CHARLES, bulletin on "Some common edible and poisonous mushrooms"	796	1–24
Pearl millet, note	793	4
Peas, fumigation with carbon disulphid, directions	799	11–12
Pelts, fox—		
demand, prices, effect of weather, etc	795	5–7
marketing, preparation and process	795	5–7, 29–31
stretching board, construction and use	795	30–31
Penicillaria, note	793	4
Pennisetum—		
glaucum, note	793	4
typhoideum, note	793	4
Pennsylvania, fur animals, laws, 1916	783	17
Pens, drainage, for sheep after dipping	798	25–26
Peppers, stuffing with mushrooms, cooking recipe	796	21
Pests—		
grain, central Oregon, control methods	800	15–16
insect, killing by carbon disulphid	799	1, 10–12, 14–21
mushroom, and how to control them, bulletin by C. H. Popence.	789	1–16
Philippe cherry, description and value	776	34
Picking, cherries, methods and equipment	776	25–29
Pigs—		
age for castration	780	3
castration—		
bulletin by Frank G. Ashbrook	780	1–6
operation, direction and suggestions	780	1–5
rupture in scrotum, occurrence	780	6
treatment in castration	780	3–5
See also Hogs.		
Pillbugs, description, occurrence on mushrooms, and control	789	9–11
"Pin" cherry, grafting stock, note	776	6
Pines, use in windbreak planting, recommendations and returns	788	12, 13, 15
Pleurotus ostreatus, description, occurrence and edible value	796	7, 9
Plowing, dry lands of Oregon, suggestions	800	12
Plunger, dipping vats, description and use	798	15
Pluteus cervinus, description, occurrence and edible value	796	10–11
Poison bait—		
sowbugs in mushroom beds	789	11
use against grain pests, formulas	800	16–17

	Farmers' Bulletin No.	Page.
Poison bran—		
formula for grasshopper control	793	27
use against grasshoppers	793	27–28
Poisoning formulas for grain pests	800	16–17
Polystictus—		
mushrooms, species and descriptions	796	15–16
species, description and occurrence	796	15–16
POPENOE, C. H., bulletin on "Mushroom pests and how to control them"	789	1–16
Porcellio laevis, description, occurrence on mushrooms, and control	789	9–11
Porcupine, bounty	783	13, 19
POTTER, GEORGE W., and ADOLPH EICHHORN, bulletin on "Contagious abortion of cattle"	790	1–12
Poults, turkey—		
care and feeding	791	16–21
lice infested, treatment	791	26
Powder-post—		
beetle. *See* Lyctus beetle.		
damage, Lyctus beetles, to seasoned hardwood, bulletin by A. D. Hopkins and T. E. Snyder	778	1–20
injury, nature, and evidence of attack	778	4
Prairie-dog, bounty	783	5
Prairies, farm lands, windbreaks	788	3–15
Prices, Sea Island cotton, factors governing	787	7–8
Prince Edward Island, fur animals, laws, 1916	783	24
Propagation, cherry trees, practices	776	5
Proso—		
notes	793	3, 4
yield in different sections, comparison with foxtail millets	793	16
Pruning, cherry, at time of planting directions	776	12–13
Publications, forage crops, list	797	35
Puffball mushrooms, descriptions and value	796	17
Puffballs, descriptions and warning	796	17
Pyrethrum—		
fumigation, use against mushroom pests	789	5–6
powder, use against springtails on mushrooms	789	9
Quebec, fur animals, laws, 1916	783	24–25
Rabbit—		
jack, bounty paid	783	5, 18
wild German, bounty	783	6
Rabbits, jack, pest in Central Oregon, damage to crops and methods of destruction	800	15–16
Raccoon, bounty paid	783	5
Ragi, note	793	4
Rattlesnake, bounty	783	8, 11, 21
REESE, H. H., bulletin on "How to select a sound horse"	779	1–27
Rhizoctonia, cause of "sore-shin" in cotton	787	32
Rhode Island, fur animals, laws, 1916	783	17–18
Rice plantations, danger from anthrax, note	784	8–9
Richmond cherry, ripening season and value	776	30
Root-knot, cotton, symptoms, cause and control	787	38
Rotation, grain growing on Oregon dry lands	800	10–11
Rotations—		
crop, value of millet	793	24–25
flax growing	785	11–13
grain growing in Maryland and Virginia	786	7–8
millet, value	793	24–25
Roughage, feeding to calves, care and management	777	8–9
Roundworms, foxes, treatment	795	25
Roup, turkeys, symptoms and treatment	791	25–26
Royal Ann cherry, description and value	776	32–33

	Farmers' Bulletin No.	Page.
Russian millet, note	793	4
Rust, cotton, causes, and control	787	39–40
Rye—		
fall sowing time, method, and rate, Maryland and Virginia	786	11, 12
growing on dry lands of Oregon	800	18–20
varieties for Maryland and Virginia	786	17–18
Sage rats, grain pest in central Oregon, control measures	800	16
Salads, mushroom, cooking recipe	796	21
Sanwa millet, note	793	4
Saskatchewan, fur animals, laws, 1916	783	25
Scarifying, sweet clover seed, machine	797	20
Schmidt cherry, description and value	776	32
Sciaria, spp. occurrence on mushrooms, note	789	3
Scours, calf, causes and treatment	777	16–17
Sea Island cotton, bulletin by W. A. Orton	787	1–40
Sea Islands, principal cotton-producing islands	787	2–3
Seed—		
bed—		
grain, preparation	786	8–9
millet, preparation	793	16–17
sweet clover, preparation	797	21–22
flax, varieties, growth habits and value	785	7–8
grain—		
for Oregon dry lands, requirements, and treatment for smut	800	13–14
selection and treatment for smuts	786	9–10
huller, type for sweet clover	797	20
millet—		
bushel, legal weight, 1917	793	24
harvesting	793	19–20
harvesting and thrashing	793	19–20
treatment for control of smut	793	26
sweet clover—		
description, choice, and treatment	797	5,7,9,19–21
germination, difficulties and treatment	797	20
hulling and scarifying	797	19–21
treatment with carbon disulphid, caution	799	21
Seed-flax production, bulletin by Charles H. Clark	785	1–20
Seeding—		
flax, time and rate	785	14–16
grain, time, method, and rate for fall sowing	786	10–12
millet, date, rate, and method	793	17–19
sweet clover, rate and methods	797	22–27
Serum vaccine—		
Animal Industry Bureau, use against anthrax	784	14–15
Sobernheim's method for anthrax control	784	13–14
Setarian, glucosid in millet hay, injury to horses	793	21
Shama millet, note	793	4
SHAMEL, A. D., bulletin on "Citrus-fruit improvement: How to secure and use tree-performance records"	794	1–16
Sheep—		
dipping—		
chutes and drainage pens	798	25–26
equipment and permanent plants, construction, etc	798	23–31
feeding millet hay	793	20
injury from dipping	798	22–23
loss from ticks	798	4
management in dipping vats	798	16
susceptibility to anthrax	784	6
tick—		
nature and eradication by dipping, bulletin by Marion Imes. *See also* Tick, sheep.	798	1–31
Shipping, foxes, direction	795	23
Siberian millet, introduction and characteristics	793	12

	Farmers' Bulletin No.	Page.
Silver fox, domesticated, bulletin by Ned Dearborn	795	1–32
Skim-milk powders, feeding to calves, preparation and value	777	11
Skins—		
fox. *See* Pelts.		
skunk, bounty paid	783	6
Slaughterhouse, offal, feed for hogs danger from tuberculosis	781	8
Slug, garden, description, occurrence on mushrooms, and control	789	11–12
Small grains. *See* Barley; Emmer; Oats; Rye; Wheat.		
Smut, millet, treatment	793	26
Smuts, grain, treatment of seed	800	14
SNYDER, T. E., and A. D. HOPKINS, bulletin on "Powder-post damage by Lyctus beetles to seasoned hardwood"	778	1–20
Soda, nitrate, use on millet	793	25
Soil—		
protection by windbreak, advantages	788	5–7
Sea Island cotton, requirements	787	3–4
Soils, worn-out, value of sweet clover for	797	13
Sore shin, cotton disease, description, cause, and prevention	787	32–33
Soup, cream of mushroom, recipe	796	21
Sour-cherry trees, pruning methods	776	17–19, 20–21
South Carolina, fur animals, laws, 1916	783	18
South Dakota, fur animals, laws, 1916	783	18
Southern Lyctus, description and distribution	778	13–14
Sowbugs, description, occurrence on mushrooms and control	789	9–11
Soy beans—		
growing in mixtures for dairy cows, practices	793	21–23
mixtures with sweet clover	797	33
Spanish cherry, description and value	776	33
Spavin, horse, symptoms	779	22
Spelt, varieties for Maryland and Virginia	786	16–17
Spraying, cherry trees, suggestions	776	24–25
Springtails, description, occurrence on mushrooms, and control	789	8–9
Squirrel, bounty	783	20
Stable—		
disinfection, control of abortion disease	790	10–11
vices, horses, corrective measures	779	3–5
Stanchions, calf, construction	777	13–14
STANTON, T. R., bulletin on "Fall-sown grains in Maryland and Virginia"	786	1–24
Sterilizer, compost box, description	789	5
Storage, flax for seed	785	19
Stretching board, casing fox skins, construction and use	795	30–31
Sudan grass—		
catch crop, comparison with millet	793	5
yields in different sections, comparison with foxtail millets	793	16
Sweeny, symptoms, cause	779	12–13
Sweet clover—		
growing the crop, bulletin by H. S. Coe	797	1–34
See also Clover, sweet.		
Sweet-cherry trees, pruning, methods	776	19–23
"Tall millet," note	793	4
Tanks, cooking and settling, for making lime-sulphur dips	798	21–22
Tanneries, anthrax transmission	784	8–9
Tartarian cherry, description and value	776	32
Tennessee, fur animals, laws, 1916	783	18
Texas—		
fever, diagnosis, distinction from anthrax	784	9
fur animals, laws, 1916	783	18
millet, note	793	4
THOMPSON, C. W., bulletin on "How the Federal farm loan act benefits the farmer"	792	1–12
THOMSON, E. H., bulletin on "The use of a diary for farm accounts"	782	1–19

	Farmer's Bulletin No.	Page.
Thrashing—		
flax	785	18–19
grain, Maryland and Virginia	786	14
Tick, sheep—		
description, life history and control by dipping, bulletin by Marion Imes	798	1–31
occurrence, prevalence, injury to sheep and wool	798	3–4
Ticks—		
detection in flock of sheep	798	10–11
sheep, eradication by dipping, principles and directions	798	11–23
spread and infestation of premises	798	9–10
Tobacco—		
fumigation, use against mushroom pests	789	5
growing after sweet clover, note	797	3
Tomatoes, baking with mushrooms, recipe	796	21
Trapping—		
fur animals, open seasons, 1916, by States and Provinces	783	27–28
legislation, 1916	783	2–3
Trappers, North American, earnings, 1915	783	1
Trees, species suited to various localities for windbreaks	788	11–13
Trichloma equestre, description, occurrence, and edible value	796	9, 12
Tuberculin test, hogs, nature and operation	781	11
Tuberculosis—		
hogs—		
bulletin by John R. Mohler and Henry J. Washburn	781	1–20
prevalence and economic importance	781	3–5
preventive measures	781	15–19
lesions in hogs, appearance and occurrence	781	12–15
Turkestan millet, introduction and characteristics	793	13–14
Turkey—		
day, practices in Eastern States	791	22
hens—		
breeding, feeding, and management	791	9–16
laying and nesting habits	791	10–12
industry in the United States	791	3–4
nesting habits	791	12
raising, bulletin by Andrew S. Weiant	791	1–27
Turkeys—		
diseases, symptoms, and treatment	791	25–26
fattening and preparation for market	791	21–24
lice infested, treatment	791	26
prices, principal markets	791	4
raising, profits	791	4
varieties, descriptions, and value	791	4–8
weights of different varieties	791	6
wild, varieties, descriptions, and occurrence	791	4–5
Tyroglyphus lintneri, description, occurrence on mushrooms and control	789	6–8
Ustilago crameri, smut of foxtail millet, control treatment	793	26
Utah, fur animals, laws, 1916	783	18–19
Vaccination, animals, against anthrax	784	11–15
Vaccine, anthrax, dangerous properties, care, etc	784	15
Vaccines, preventive, use against anthrax	784	11–15
Vat—		
cement, sheep dipping, construction	798	30–31
wooden, sheep dipping, construction	798	30
Vats, dipping, for sheep, portable, and permanent	798	23–24, 26–31
Velvet Lyctus, description and distribution	778	14–15
Vermont, fur animals, laws, 1916	783	19
VINALL, H. N., bulletin on "Foxtail millet: Its culture and utilization in the United States"	793	1–28

	Farmers' Bulletin No.	Page.
Virginia—		
fall-sown grain, and in Maryland, bulletin by T. R. Stanton	786	1–24
fur animals, laws, 1916	783	19–20
grain acreage, by varieties, 1911–1915	786	3
grain growing, fall sowing, time, rate, and varieties	786	24
grain production, yields of various kinds, 1906–1915	786	21–22
Vixen, fox, habits and treatment	795	20–21
WASHBURN, HENRY J.—		
and JOHN R. MOHLER, bulletin on "Tuberculosis of hogs"	781	1–20
bulletin on "Anthrax or charbon"	784	1–16
Washington, fur animals, laws, 1916	783	20
Weasel, bounty	783	6, 17
Weeds, destruction by millet crop	793	7
Weeping merulius mushroom, description	796	16
WEIANT, ANDREW S., bulletin on "Turkey raising"	791	1–27
Weighing, citrus fruit in orchard, management	794	6–12
West Virginia—		
fur animals, laws, 1916	783	20
grain—		
acreage, by varieties, 1911–1915	786	3
production, yields of various kinds, 1905–1915	786	21
Western Lyctus, description and distribution	778	15
Wheat—		
fall sowing, time, method, and rate, Maryland and Virginia	786	10, 11, 12
growing on dry lands of Oregon	800	17–18
value, by acre, comparison with flax, oats, and barley	785	6
varieties for Maryland, Virginia, and Delaware	786	15–16
White—		
Holland turkey, description, weight, and value	791	5, 6
scours, calf disease, nature and treatment	777	16–17
Smyrna barley, characteristics and value for dry lands of Oregon	800	21
Wildcat, bounty paid, by States and Provinces	783	5, 7, 10–13, 17, 21, 23
Willow, use in windbreak planting, recommendations and returns	788	13, 14
Wilt, cotton, symptoms, cause and control	787	35–38
Wind movements, effect of windbreaks	788	3–4
Windbreak—		
contour, requirements for best service	788	4
effect on—		
crop yields, determination	788	9–11
temperature, importance in corn growing, etc	788	7–8
farm asset, bulletin by Carlos G. Bates	788	1–16
planting, species and location	788	11–14
protection of crops, houses, and soil	788	3–8
Windbreaks—		
composition, width, and money returns from trees	788	11–15
wood supply, yield of various species	788	14
Winds, injurious, control by windbreaks, various localities	788	3–8, 11–13
Windsor cherry, nature, ripening season and value	776	32
Wisconsin, fur animals, laws, 1916	783	20–21
Wolf, bounty paid by States and Provinces	783	4–5, 7–8, 11–13, 16–25
Woodchuck, bounty	783	7, 11, 14
Woodlice, description, occurrence on mushrooms, and control	789	9–11
Wragg cherry, nature, ripening season, and value	776	31
Wyoming, fur animals, laws, 1916	783	21
Yellow Spanish cherry, description and value	776	33
Yukon, fur animals, laws, 1916	783	25

XUM